

Tabla n° 1: Criterios de búsqueda	
Entrada de búsqueda	tipo de estudio
hepatitis C AND (neuropsychiatric OR psychiatric OR depression OR anxiety OR psychosis OR mania OR hypomania OR Bipolar)	controlled/uncontrolled clinical trial, randomized clinical trial, Meta-analysis
interferon AND (neuropsychiatric OR psychiatric OR depression OR anxiety OR psychosis OR mania OR hypomania OR Bipolar)	controlled/uncontrolled clinical trial, randomized clinical trial, Meta-analysis
hepatitis C AND interferon AND (neuropsychiatric OR psychiatric OR depression OR anxiety OR psychosis OR mania OR hypomania OR Bipolar OR quality of life)	controlled/uncontrolled clinical trial, randomized clinical trial, Meta-analysis
hepatitis C AND interferon AND treatment AND (neuropsychiatric OR psychiatric OR depression OR anxiety OR psychosis OR mania OR hypomania OR Bipolar)	controlled/uncontrolled clinical trial, randomized clinical trial, Meta-analysis
interferon AND Hepatitis C AND ((neuropsychiatric OR psychiatric OR depression OR anxiety OR psychosis OR mania OR hypomania OR Bipolar) AND quality of life))	controlled/uncontrolled clinical trial, randomized clinical trial, Meta-analysis




Tabla n° 2: Síntomas psicosomáticos en pacientes con hepatitis C	
cansancio*	86%
irritabilidad	75%
depresión	70%
agotamiento mental*	70%
olvido	65%
dificultades para dormir*	65%
disminución de la concentración	62%
dolor corporal generalizado	57%
ira	11,3%
*percepción de autoreporte con mayor relevancia	

Tabla n° 3: Prevalencia de trastornos psiquiátricos en población de infectados con el virus de la hepatitis C	
depresión	12% - 55,5%
trastorno de ansiedad	13,1% - 50%
trastorno bipolar	4%
dependencia alcohólica	35,4% - 55%
abuso de sustancias ilícitas	45% - 69,2%
indiferenciados	18,57%
metadona	25,70%
drogas intravenosas	74%
tabaquismo	44%

Tabla n°4: Trastornos psiquiátricos en pacientes infectados con hepatitis C, en tratamiento con interferón*	
depresión	16,6% - 50,9% (con un peak a las 8 y 12 semanas)
trastorno de ansiedad	16% - 25,6%
psicosis	3,30%
*interferón alfa2a pegilado y no pegilado	

TABLA N° 5: Escalas psicométricas utilizadas en el estudio de la hepatitis C	
SIGLA	NOMBRE
Métodos de screening psiquiátrico	
SCID DSM IV	Entrevista estructurada del <i>Diagnostic and Statistical Manual</i>
BPRS	Brief Psychiatric Scale
MINI	Mini-International Neuropsychiatric Interview
SCID I DSM IV	Structured Clinical Interview for DSM-IV Axis I Disorders
SCL-90-R	Self-Report Symptom Inventory 90 Items-Revised
Depresión	
MADRS	Montgomery Asberg Depression Rating Scale
CES - D	Center for Epidemiologic Studies Depression
IDD	Inventory to Diagnose Depression
HDRS	The 21-item Structured Interview Guide for the Hamilton D
BDI	The Beck Depression Inventory
ZSDS	The Zung Self-Rating Depression Scale
PMS	Profile of Mood States
Ansiedad	
HADS	Hospital Anxiety and Depression Scale
HAM - A	Hamilton Rating Scale for Anxiety
Calidad de Vida y Fatiga	
SF - 36	Short-Form 36 questions Health Survey for general health r
CLDQ	Chronic Liver Disease Questionnaire
BFI	Brief Fatigue Inventory.
MFI	Multidimensional Fatigue Inventory
Cognitivas	
CANTAB	Cambridge Neuropsychological Test Automated Battery
Otras	
STAXI - 2	Anger The State–Trait Anger Expression Inventory
PNS	Positive and Negative Affect Scale
PSQI	Pittsburgh Sleep Quality Index

Tabla n° 6: Distribución de síntomas en pacientes con hepatitis C en tratamiento con interferón

ánimo depresivo	35% - 64%
ansiedad	25,6%
ira/hostilidad	24,5% - 75%
manía	10%
hipomanía/irritable	50%
psicosis	3,3%
anorexia	63%
alteración de la concentración	50%
somatización	8,2%
fobia social	4,5%
pensamiento paranoide	3,1%